



## Phytochemical Profiling and Evaluation of Antioxidant and Antidiabetic Activity of Methanol Extract of Spinach (*Spinacia Oleracea* L.) Leaves

Ohidul Islam<sup>1</sup>, A.K. Azad<sup>2</sup>, M. Mustafezur Rahman<sup>2</sup>, A. Khorshed Alam<sup>1</sup>, M. Khairuzzaman<sup>1</sup>, Jannatun Ferdous<sup>1</sup>, Manirul Islam<sup>1</sup>, Sharif M. Shaheen<sup>\*2</sup>

<sup>1</sup>Department of Pharmacy, Bangladesh University, Dhaka, Bangladesh

<sup>2</sup>Department of Pharmacy, Daffodil International University, Dhaka, Bangladesh

### Abstract

The purpose of this study was to investigate the presence of major phytochemicals and the anti-diabetic and antioxidant effects of the methanolic extract of the plant *Spinacia oleracea* L. leaves. Phytochemical analysis of *Spinacia oleracea* L. displayed the presence of alkaloids, saponins and tannin types of compounds. The anti-oxidative effect was evaluated using DPPH free radical scavenging activity method. The methanolic extract of *Spinacia oleracea* L. leaves is found to have no antioxidant activity. The IC<sub>50</sub> of the extraction is 211081.58 µg/ml. In case of anti-diabetic activity test good anti-diabetic activity was observed. Mice treated with extract Group (250 mg/kg) showed decrease (from 21.7 mM ± SEM to 14.1 mM ± SEM) (p<0.05) in blood glucose concentration at 120 min and extract group (500 mg/kg) showed decrease (from 24.9 mM ± SEM to 14.3 mM ± SEM) (p<0.05) at 120 min compared with Standard Group (from 25.6 mM ± SEM to 11.2 mM ± SEM) at 120 min.

**Keywords:** Phytochemical Profiling, Antioxidant, Antidiabetic, DPPH, Alloxan, Carrageenan

### Corresponding author: Sharif Mohammad Shaheen

Department of Pharmacy, Daffodil International University, Dhaka, Bangladesh.

**E-mail:** [smsshaheen2001@yahoo.com](mailto:smsshaheen2001@yahoo.com); [sharif.ph@diu.edu.bd](mailto:sharif.ph@diu.edu.bd)

**Citation:** Sharif Mohammad Shaheen, et al (2017). Phytochemical Profiling and Evaluation of Antioxidant and Antidiabetic Activity of Methanol Extract of Spinach (*Spinacia Oleracea* L.) leaves... Int J Pharm Sci & Scient Res. 3:8, 24-27

**Copyright:** ©2018 Sharif Mohasmmad Shaheen et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Received:** December 26, 2017

**Accepted:** January 06, 2018

**Published:** March 07, 2018

### Introduction

Considerable portion of current diseases are caused due to the 'oxidative stress' which results in enormous amount of free radicals, causing tumor, atherosclerosis and cardiovascular illnesses (Braca et al., 1999). Cells of the human body ensure themselves against harm caused by free radicals by catalysts such as ascorbic acid, tocopherol and glutathione (Braca et al., 2016). Cell reinforcement supplements are imperative to battle oxidative harm. That is why much consideration has been taken towards the improvement of ethnomedicine with solid cell reinforcement properties with low cytotoxic effects.

Diabetes mellitus (DM) is a metabolic and endocrine disorder which currently affects more than 100 million people all over the world and the number of affected people is increasing due to aging, increasing prevalence of obesity and physical inactivity (Nair et al., 2006; Sarah et al., 2004; Safdar et al., 2004; Umara et al., 2010). According to recent studies, approximately by the year 2030, 438 million people all over the world are expected to have diabetes (Mojekwu et al., 2011; Noor et al., 2008; Rehman et al., 2011). The worldwide cost to control diabetes and associated complications exceeds \$100 billion per year and complications are far less common and less severe in people who have well controlled blood sugar levels (Chattopadhyay, 1999; Sokeng et al., 2001). The treatment of diabetes with synthetic drugs is generally not preferred because of its high cost and side effects, for this reason, it is necessary to develop alternative medicines of plant based origins with anti-diabetic properties (Emmanuel et al., 2010; Tanko et al., 2008).

*Spinacia oleracea* L. (Spinach) is a leafy green vegetable that came originally from southwestern Asia and is now grown in most parts of the world. Its leaves, which are broad and smooth and about ten inches long, Spinach, especially raw, is a very good source of folic acid, Spinach leaves are rich in vitamin C and E, which are antioxidant.

These are supposed to lower risks of heart disease, stroke and cancer. The high amount of vitamin A in spinach may protect against eye degeneration. The potassium helps prevent and regulate high blood pressure. The plant is carminative and laxative. In experiments it has been shown to have hypo-glycemic properties. The aim of this study was to determine the presence of major phytochemicals in methanolic extract of *Spinacia oleracea* L. leaves and to investigate its anti-oxidative and anti-diabetic effects.

## Materials and methods

### Plant Materials

*Spinacia oleracea* L. leaves were collected from Mahammadpur, Townhall Kacha Bazar, Dhaka and the plant authentically was confirmed from the Bangladesh National Herbarium,

### Drying and Grinding

The collected plants were separated from undesirable materials or plants or plant parts. They were dried in the sun for one week after cutting into small pieces. The plant parts were ground into coarse powder with the help of a suitable grinder. The powder was stored in an airtight container and kept in a cool, dark and dry place until analysis commenced.

### Preparation of Plant Extract

About 300 gm of powdered sample was taken in a clean, flat-bottomed glass container and soaked in 1500 ml of 90% methanol. The container with its contents was sealed and kept for a period of 10 days accompanying occasional shaking and stirring. The whole mixture then underwent a coarse filtration by a piece of clean, white cotton material. Then it was filtered through Whatman filter paper. The filtrate was kept in an open space to evaporate the solvent thus crude extract was obtained. Fine powders of the flowering plant of *Spinacia oleracea* L. leaves are dissolved in 90% methanol and then evaporation of the solvent.

### Phytochemical Screening

Phytochemical studies of methanolic extract of plant material extract was carried out for preliminary chemical investigation for the direction of practical pharmacognosy text book (Trease and Evans, 1983; Mohammed Ali, 2012; Abdul Ghani, 2005).

**Antioxidant tests** (Proctor, 1989; Hennekens et al., 1994; Clarkson, 1995)

Stock solution of the plant extract was prepared in methanol (10mg/ml) from which a serial dilution was carried out. At first 6 volumetric flasks are taken to make 6 different types of concentration 1, 5, 10, 50, 100 and 500 µg/ml. Test tubes and volumetric flasks are capped with foil paper. In 6 volumetric flasks serial dilution of extract is done and marked them respectively. 2ml of sample from each concentration and 2 ml of 0.004% DPPH solution is taken with the help of pipette in 6 test tubes respectively.

2ml of sample from each concentration and 2 ml of 0.004% DPPH solution is taken with the help of pipette in 6 test tubes respectively. Then solution is kept in dark place for 30 minutes with rapping each test tube with foil paper. In another test tube 2ml 0.004% DPPH & 2ml methanol is taken to prepare blank solution. Then absorbance is taken by UV Spectroscopy. The percent of inhibition is calculated by using following formula

$$\% \text{ of inhibition} = \frac{\text{Blank absorbance} - \text{Solution absorbance}}{\text{Blank absorbance}} \times 100$$

### Drugs and chemicals

Carrageenan was purchased from Otto chemicals, India. The standard drug Diclofenac-Na was purchased from Square Pharmaceuticals Limited of Bangladesh. Acetic acid, methanol and other chemicals supplied from laboratory of Bangladesh University were analytical grade.

### Experimental animals

Eight week-old Swiss albino mice (27-30g) purchased from Jahangirnagar University, Dhaka, Bangladesh and were housed in animal cages under standard environmental conditions (22-25°C, humidity 60-70%, 12 hr light: 12 hr dark cycle). The mice were fed with standard pellet diet taken from Jahangirnagar University Dhaka. The animals used in this study were cared in accordance with the guidelines on animal experimentation of our institute.

## Method for Evaluation of Hypoglycemic Activity

### Oral Glucose Tolerance Test (OGTT) in diabetic mice

After fasting 16hr, diabetes was induced into mice by intra-peritoneal injection (i.p.) of alloxan monohydrate (90 mg/kg) dissolved in saline. After 48hrs, plasma glucose levels were measured by glucometer (Tyson, Taiwan) using a blood sample from tail-vein of mice. Mice with blood sugar higher than 11.5 mmol/l were considered as diabetic. All the mice were divided into 4 groups, each group containing 5 mice. The divided groups are NC (normal control), DC (diabetic control), STD (diabetic mice receiving Metformin), ME (diabetic mice receiving methanolic extract). The mice were fasted overnight and next day blood samples were taken from all groups of animals to estimate fasting blood glucose level (0 min). All mice received 1gm/kg glucose. Without delay extract and were given per oral and three more blood samples were collected at 30, 90 and 120 minutes intervals and blood glucose level was estimated in all the experiments by using glucometer (Hossain et al., 2011).

## Result and discussion

### Phytochemical Screening

Phytochemical screening of Methanolic Extract of *Spinacia oleracea* L. leaf is displayed in Table-1

Chemical Groups	Methanolic Extract of <i>Spinacia oleracea</i> L. leaf
Saponin	+
Glycoside	-
Flavonoids	-
Tannin	+
Alkaloids	+

**Table 1:** Results of Phytochemical Screening


Note: (+) = Indicates the presence and (-) = Indicates the absence of the tested group.

**Result of Anti-oxidants test**

DPPH scavenging assay was used to determine the antioxidant activity

Conc. (µg/ml)	Absorbance (nm)		% of Inhibition		
	Blank	Ascorbic Acid	<i>Spinacia oleracea L.</i>	Ascorbic Acid	<i>Spinacia oleracea L.</i>
1	0.902	0.863	0.872	4.3	3.3
5		0.780	0.864	13.6	4.2
10		0.509	0.849	43.6	5.9
50		0.047	0.834	94.8	7.5
100		0.034	0.797	96.2	11.6
500		0.023	0.598	97.5	33.7

**Table 2:** % Inhibition of Ascorbic acid and *Spinacia oleracea L.*



**Fig 1.** Anti-oxidant activity of Ascorbic acid and *Spinacia oleracea L.*

Test Samples	Regression line	R2	IC <sub>50</sub> µg/ml
Ascorbic Acid	$y = 18.114\ln(x) + 2.0487$	$R^2 = 0.8726$	14.15
<i>Spinacia oleracea L.</i>	$y = 4.2575\ln(x) - 2.196$	$R^2 = 0.6898$	211081.58

**Table 3:** IC<sub>50</sub> values of the extracts of Ascorbic Acid and *Spinacia oleracea*

The IC<sub>50</sub> of *Spinacia oleracea L.* is 211081.58µg/ml, whereas IC<sub>50</sub> of Ascorbic Acid is 14.15µg/ml.

**Results of Hypoglycemic Activity**

Animal Group	0 min	30 min	60 min	90 min	120 min
Control Group	18.5±1.09	21.3±0.53	19.7±0.44	16.6±0.66	15.2±0.69
Standard Group	25.6±0.44***	28.5±0.38***	16.3±0.59	12.8±0.69**	11.2±0.74***
Extract Group (250 mg/kg)	21.7±0.33*	27.4±0.36***	21.8±0.49*	18.6±0.52	14.1±0.41
Extract Group (500 mg/kg)	24.9±0.51***	27.8±0.38***	21.2±0.64	17.9±0.53	14.3±0.31

**Table 4:** Oral Glucose Tolerance Test (OGTT) of *Spinacia oleracea L.* leaf extract in alloxan-induced diabetic (mM/L) in mice

Experimental data were presented as mean ± SEM. By using the Dunnett test significant differences (\*p<0.05, \*\*p<0.01, \*\*\*p<0.001) between the means were determined compare to control group where n=04. For statistical evaluation IBM-SPSS software version 20 was utilized.

## Discussion

This study demonstrated the findings of Phytochemical studies, antioxidant and anti-diabetic activity of methanolic extract of *Spinacia oleracea* L. leaves by several in vivo and in vitro method. Phytochemical observations have revealed the presence of several phytochemicals including alkaloids, saponins, and tannins.

The reducing power of a compound may act as a momentous indicator of its potential antioxidant activity (Pal R et al., 2011). Samples with elevated reducing power are better capable to donate the electron and free radical from stable substance by accepting the donated electrons, resulting in the termination of radical chain reaction (Deori M et al., 2014).

The experiment showed with  $IC_{50}$  of the extract is 211081.58 $\mu$ g/ml, whereas  $IC_{50}$  of Ascorbic acid is 14.15 $\mu$ g/ml. When the  $IC_{50}$  value of extract compared with the standard ascorbic acid it seems a large value. Which unfortunately poses that, the extract offers no antioxidant activity (Wasim M et al., 2015). But, the extract presents a great positive effect on mice with alloxan induced disturbance in glucose tolerance.

Alloxan is a popular diabetogenic agent hydrophilic in nature and chemically unstable pyrimidine derivative, which harms pancreatic  $\beta$ -cells because it can generate toxic free oxygen radicals during redox cycling in the presence of reducing agents such as glutathione and cysteine (Wasim M et al., 2015).

Experimental data were presented as mean  $\pm$  SEM. By using the Dunnett test significant differences ( $*p < 0.05$ ,  $**p < 0.01$ ,  $***p < 0.001$ ) between the means were determined compare to control group where  $n=04$ . For statistical evaluation IBM-SPSS software version 20 was utilized.

## Conclusion

Phytochemical analysis of *Spinacia oleracea* L. displayed the presence of alkaloid, saponin and tannin types of compounds. The results stated above showed that the methanolic extract of *Spinacia oleracea* L. possessed no antioxidant effect and very good anti-diabetic properties. However, this can't be confirmed without further higher and specific tests. So, further researches should be conducted to get information about these activities.

## Conflict of interest

The authors declare that they have no conflict of interest.

## Acknowledgement

The authors are grateful to Bangladesh University for providing the space for research and necessary lab equipments.

## References

1. Abdul Ghani. Practical Phytochemistry, 1st ed., Dhaka: Parash Publishers; 2005: 412-418.
2. Braca A, Sortino C, Politi M, Morelli I, Mendez J. Antioxidant activity of flavonoids from *Chattopadhyay RR*. Possible mechanism of antihyperglycemic effect of *Azadirachta indica* leaf extract. *J Ethnopharmacol* 1999; 67(3):373-376
3. Clarkson PM. Antioxidants and Physical Performance. *Crit Rev Food Sci Nutr*, 1995; 35(1&2): 131-141.
4. Emmanuel S, Rani S, Sreekanth R. Antidiabetic activity of *Cassia occidentalis* in streptozotocin-induced diabetic rats: a dose dependent study. *IJPBS*. 2010; 1: 1-12.
5. Hennekens CH, Buring JE, Peto R. Antioxidant Vitamins Benefits Not Yet Proved. *N Engl J Med*. 1994; 330(15): 1080 – 1081.

6. Hossain MS, Asadujjaman M, Khan MRI, Ahmed M, and Islam A, Antidiabetic and glycogenesis effects of different fractions of methanolic extract of *Momordica charantia* (Linn.) in alloxan induced diabetic rats. *Int J Pharm Sci Res*. 2011; 2(2): 404-412.
7. Meetal Deori, Dulal Chandra Boruah, Dipali Devi, Rajlakshmi Devi; Antioxidant and antigenotoxic effects of pupae of the muga silkworm *Antheraea assamensis*. *Food Bioscience* 5, 2014: 108-114.
8. Mohammad Wasim, Kuldeep Singh, MK Mishra; Antidiabetic, antihyperlipidemic and histopathological studies of aqueous and ethanol extracts of leaves of *Scirpus grossus* in alloxan induced diabetic rats. *International Journal of Pharmaceutical Sciences Letters* 2015, 5 (6): 627-631.
9. Mohammed Ali. Textbook of Pharmacognosy, 2nd ed., Delhi: CBS Publishers & Distributors Pvt. Ltd.; 2012: 141-175.
10. Mojekwu TO, Yama OE, Ojokuku SA, Oyebadejo SA. Hypo glycaemic effects of aqueous extract of *Aframomum melegueta* leaf on alloxan-induced diabetic male albino rats. *Pac J Med Sci*. 2011; 8(1):28-36
11. Nair SA, Shylesh BS, Gopakumar B, Subramoniam A. Antidiabetes and hypoglycaemic properties of *Hemionitis arifolia* (Burm.) Moore in rats. *J Ethnopharmacol*. 2006; 106(2): 192-197.
12. Noor A, Gunasekaran S, Soosai A, Minicab, Vijayalakshmi MA. Antidiabetic activity of Aloe vera and histology of organs in streptozotocin induced diabetic rats. *Current science* 2008; 94:1070-1076.
13. Proctor PH. Free radicals and human disease, In: Miquel J; Weber H., eds. *Handbook of Free Radicals and Antioxidants in biomedicine*, 1st ed. Florida: CRC Press; 1989: 209-221.
14. Ranju Pal, Kundlik Girhepunje, Nidhi Shrivastav, Mohammed Misbah Hussain and Thirumoorthy N; Antioxidant and free radical scavenging activity of ethanolic extract of *Morinda citrifolia*. *Annals of Biological Research*, 2011, 2 (1): 127-131.
15. Rehman SU, Jafri SA, Hassan S, Ishtiaq N, Muhammad N. Study on antidiabetic effect of Aloe vera extract on alloxan induced diabetic rats. *LARCJI*. 2011; 2(1): 29-32.
16. Safdar M, Khan A, Khan MMA, Siddique M. Effect of Various Doses of Cinnamon on Blood Glucose in Diabetic Individuals. *PJN*. 2004; 3(5):268-272.
17. Sarah W, Anders G, Sicree R, King H. Global Prevalence of Diabetes: epidemiology/health services/psychosocial research. *Diabetes Care* 2004; 27(5): 1047-53.
18. Sokeng DS, Lontsi D, Moundipa PF, Jatsa HB, Watcho P, Kamtchoung P. Hypoglycemic effect of *Anacardium occidentale* L Aqueous extract in normal and on streptozotocin-induced diabetic rats. *Diabetes Res*. 2001; 36: 01-09.
19. Tanko Y, Yerima M, Mahdi MA, Yaro AH, Musa KY, Mohammed A. Hypoglycemic Activity of Methanolic Stem Bark of *Adansonia digitata* extract on blood glucose levels of Streptozotocin-induced Diabetic wistar rats. *Int. j. appl. res. nat. prod*. 2008; 1(2):32-36.
20. Trease GE, Evans WC. Textbook of Pharmacognosy, 12th ed., London: Bailliere Tindall and Company Publisher; 1983: 343-383.
21. Umara A, Qamar U, Bala Y, Bashar Bello MS. Anti-hyperglycemic activity of the leaves of *Tetracerascandens* (Dilleniaceae) in alloxan induced diabetic rats. *J Ethnopharmacol*. 2010; 131:140-145.
22. Valko M, Jomova K, Rhodes CJ, Kuca K, Musilek K. Redox- and non-redoxmetal-induced formation of free radicals and their role in human disease. *Arch Toxicol*. 2016;90(1):1-37.