

## Alterations in the Leucocytes and Serum Biochemistry in grey mullet ( Mugil cephalus L , ) Fingerlings exposed to Sub Lethal Doses of Lead for different Exposure periods

Hussien. M. EL-Shafei\*

General Authority for Fish Resource Development, Egypt

**Corresponding Author:** Hussien. M. EL-Shafei, General Authority for Fish Resource Development, Egypt.

E-mail: [hussienelshafei50@yahoo.com](mailto:hussienelshafei50@yahoo.com)

**Citation:** Hussien. M. EL-Shafei (2017), Alterations in the Leucocytes and Serum Biochemistry in Grey Mullet (Mugil cephalus L.) Fingerlings exposed to Sub Lethal Doses of Lead for different Exposure periods. Int J Biotech & Bioeng. 3:6, 165-170.

DOI: [10.25141/2475-3432-2017-6.0159](https://doi.org/10.25141/2475-3432-2017-6.0159)

**Copyright:** ©2017 Hussien. M. EL-Shafei. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Received:** April 20, 2017; **Accepted:** May 03, 2017; **Published:** July 05, 2017

### Abstract:

Grey Mullet *Mugil cephalus* L, fingerlings were chronically exposed to sub lethal concentrations of lead(Pb) (0.1 and 0.4 mg/L for twenty eight. The changes in the leucocytes and serum glucose, protein, and total cholesterol of the fish were determined every seven days in a renewable static bioassay system. At the end of the study, these parameters were significantly ( $p < 0.05$ ) elevated in the Pb - exposed groups when compared with the control group. Showing a pronounced leucytosis in the Pb-exposed fish. The magnitude of increase was influenced by increasing of exposure period and Pb concentration. The Pb exposed fish were significantly ( $p < 0.05$ ) hyperglycemic and hypercholesteremic. The serum glucose levels on the 7th day were  $26.50 \pm 2.12$  g/dl and  $30.50 \pm 0.70$  g/dl in the fish exposed to 0.1 and 0.4 mg Pb /L respectively. On the 28th day, the serum glucose concentrations were  $52.50 \pm 2.12$  and  $70.00 \pm 2.83$  g Pb /dl in the groups exposed to 0.1 and 0.4 mg Pb/L, respectively. The cholesterol concentration increased from  $113.5 \pm 3.53$  mg Pb /dl on day 7 to  $208.0 \pm 1.80$  mg/dl on day 28 in the group exposed to 0.1mg Pb /L. When the fish was exposed to 0.4mg Pb /L lead acetate, the cholesterol concentration increased from  $131.5 \pm 3.54$  in the first week to  $288 \pm 5.19$ mg Pb /dl on 28th day of the study. The serum protein concentration was also significantly ( $p < 0.05$ ) increased in the Pb – exposed groups when compared with the control group, it increased from  $4.04 \pm 0.06$ mg Pb /dl on day 7 to  $5.30 \pm 0.05$  mg/dl on day 28 in the fish exposed to 0.1mgPb /L. When the fish treated with 0.4mg Pb /L, the serum protein increased from  $4.45 \pm 0.37$  mg/dl on day 7 to  $6.18 \pm 0.19$  mg Pb /dl on day 28, respectively. These changes are indicative of stress imposed on the fish by lead and could be used as indices of lead poisoning

**Keywords:** Lead, Hyperglycemia, Cholesterol, Protein, Leukocyte

### Introduction:

Lead(Pb) is one of the most widely used heavy metals that has wide applications in such products as storage batteries (lead accumulator), electric cable sheaths, alloys, pesticides, paints, petrol and rubber products among other uses [1]. In water bodies, Pb forms complexes with sediments or organic materials [2]. and in the process enters the food chain. The problem posed by Pb in the aquatic system is complex due to its non-degradability and interactions with other materials to form complexes that may potentiate their toxic effects[3] Lead deposits in various

fish organs: liver, kidneys and spleen, but also digestive tract and gills [4]. Accumulation of lead in different fish species has been determined in several works[5], leading to disorders in fish body. When *C. batrachus* exposed to 5 ppm of lead nitrate for 150 days, it exhibited marked inhibition of gonadal growth and showed decrease in cholesterol and lipid levels in brain, testis and ovary whereas the liver showed an elevation of both [6]. [7] observed a very high number of red let cells (RCs) in the epidermis of common carp and rainbow trout kept in lead polluted water. Hepatocyte vacuolization, hepatic cirrhosis, necrosis, shrinkage, parenchyma degeneration, nuclear psychosis and increase of sinusoidal spaces

were the distinct changes observed in the liver of lead-exposed fish [8]. The characteristic symptoms of chronic lead toxicity include changes in the blood parameters with severe damage to erythrocytes and leucocytes and damage in the nervous system [9]. Low levels of Pb pollution could cause some adverse effects on fish health and reproduction [10]. Also, lead was found to impair the embryonic and larval development of fish species [11]. [12] monitored the effect of lead on the Chinese sturgeon, *Acipenser sinensis*. They observed deformities as body (spinal) curvatures. The authors also reported reduced ability of locomotion and foraging by deformed juveniles. The purpose of the study was therefore to investigate the effect of sub lethal lead on the leucocyte and some biochemical parameters in grey mullet *Mugil cephalus*, L- with a view to using them as biomarkers of lead toxicity.

### Materials and methods:

Collection and handling of experimental of fish.

The fish fingerlings were bought from Fish Farms at Manzalla lake Damietta, and transported to a wet laboratory in two 50 L plastic fish transport containers during march 2015. Once in the wet laboratory, fish was introduced into 200 L plastic container for acclimatization for two week. The water was continually aerated to ensure that the dissolved oxygen level remained above 6 mg/L and fish were fed on a 32% crude protein diet at 3% body weight daily at 10.0 h. The laboratory water was analyzed for different physico-chemical parameters (APHA, 2010) and lead. No fish mortality was recorded during acclimatization. The LC50 value for lead was statistically determined [13]

Experimental design and in vivo studies

One hundred and twenty fingerlings of *Mugil Cephalus* L, (mean length of 12.00±2.25 cm and average weight of 35±3.20g) were used for this study. They were divided into three groups of forty fingerlings per each. Each group was further divided into two replicates, twenty fish per replicate. Fish were exposed to 0.1 mg Pb /L and the second group was exposed to 0.4 mg Pb /L.

### Result:

Water quality parameters, cations, anions, and background metals in acclimation. were presented in table(1)

Parameter	Acclimation water
Temperature (°C)	25±1°C
pH	8.1±0.2
Dissolved oxygen (mg/l)	8.2-8.9
Total Hardness (mg/l as CaCO <sub>3</sub> )	106.2
Total alkalinity(mg/l as CaCO <sub>3</sub> )	42.8
Total dissolved solids(mg/l)	173
Sodium (mg/l)	5.0
Calcium (mg/l)	31.0
l)Potasium(mg)	0.6
Magnesium (mg/l)	6.1
Cl- (mg/l)	10.5
NH <sub>3</sub> (mg/l)	0.034
SO <sub>4</sub> (mg/l)	13.0
PO <sub>4</sub> (mg/l)	0.03
Copper(µg/l)	0.63
lead (µg/l)	0.015
Zinc (µg/l)	0.35

**Table (1):** Water quality parameters, cations, anions, and background metals in acclimation

The third group was exposed to tap water only and it served as control group. The water in the experiment was changed every day to maintain the same Pb concentrations. Three fish from each replicate experiment was killed every seven days for twenty eight days for the analysis.

### Blood Collection

The blood of the fish was collected every seven days through both cardiac puncture [14]. The blood was collected into three different vials. One of the vials containing fluoride oxalate was used to collect blood used for glucose determination. The other two vials were without anticoagulant and were used to collect the blood for the biochemical test, and thin blood smear.

### White Blood Cell Count (WBCC):

The leucocyte count was done using the Neubauer microscopic counter after diluting the blood with Turk's dilution fluid. The differential white blood cell count was done by preparing a thin blood smear and staining same with Geimsa Romanosky stain [15]. The stained blood was left for 25 minutes for the Geimsa stain to act on it. Thereafter the stained blood was flooded with distilled water and rocked gently to evenly mix the distilled water and stain. The stained blood was washed with water and allowed to dry. The slide was viewed with a binocular microscope to identify the leukocyte species which was calculated as a percentage.

### Biochemical analyses:

The blood serum was obtained by centrifuging the blood sample at 5000 rpm for five minutes and the protein in the samples was spectrophotometrically determined using Biuret method [16] at 540 nm. The blood glucose was determined by [17] method and the cholesterol concentration was determined by the method of [18].

### Statistical Analysis:

The data was statistically analyzed using two way analysis of variance (ANOVA) followed by LSD post hoc test at 95% confidence interval [19].

### Leukocyte Response

The changes in the total leucocytes and differential white blood cell count Pb- exposed to Fish are presented in (Table 2)

Leucocyte species	Concentration Pb(mg/L)	Duration( days)			
		7	14	21	28
Lymphocyte	Control	81.00±1.41 <sub>a1</sub>	80.50±3.54 <sub>a1</sub>	80.00±1.41 <sub>a1</sub>	85.50±0.71 <sub>a1</sub>
	0.10	83.50±2.12 <sub>b1</sub>	84.50±0.71 <sub>b2</sub>	85.50±2.12 <sub>b3</sub>	87.50±3.54 <sub>b4</sub>
	0.40	83.50±2.12 <sub>b1</sub>	86.00±2.83 <sub>b2</sub>	87.00±1.41 <sub>c3</sub>	87.00±1.41 <sub>b4</sub>
Monocyte (%)	Control	2.50±0.71 <sub>a1</sub>	2.30±1.41 <sub>a1</sub>	2.40±0.71 <sub>a1</sub>	2.50±0.71 <sub>a1</sub>
	0.10	1.00±1.41 <sub>b1</sub>	5.50±1.41 <sub>b2</sub>	6.00±1.41 <sub>b2</sub>	4.00±0.00 <sub>b2</sub>
	0.40	0.50±0.71 <sub>c1</sub>	6.00±2.83 <sub>c2</sub>	5.00±1.41 <sub>c3</sub>	5.50±1.41 <sub>c3</sub>
Neutrophil (%)	Control	1.50±0.71 <sub>a1</sub>	1.50±0.71 <sub>a1</sub>	1.50±0.71 <sub>a1</sub>	1.30±0.71 <sub>a1</sub>
	0.10	1.00±1.41 <sub>b1</sub>	2.00±1.41 <sub>b2</sub>	1.50±0.71 <sub>a3</sub>	1.50±0.71 <sub>a3</sub>
	0.40	0.50±0.71 <sub>c1</sub>	1.50±0.71 <sub>c2</sub>	0.50±0.71 <sub>b1</sub>	2.00±1.41 <sub>b3</sub>
Eosinphil (%)	Control	0.50±0.71 <sub>a1</sub>	0.50±0.71 <sub>a1</sub>	0.60±0.71 <sub>a1</sub>	0.50±0.71 <sub>a1</sub>
	0.10	0.50±0.71 <sub>a1</sub>	1.00±1.41 <sub>b2</sub>	0.50±0.71 <sub>a1</sub>	-
	0.40	-	0.50±0.71 <sub>a1</sub>	0.50±0.71 <sub>a1</sub>	0.50±0.71 <sub>a1</sub>
Basophil (%)	Control	10.50±0.71 <sub>a1</sub>	10.50±2.12 <sub>a1</sub>	9.80±2.12 <sub>a1</sub>	10.50±2.12 <sub>a1</sub>
	0.10	15.50±0.71 <sub>a1</sub>	5.50±0.71 <sub>b2</sub>	8.50±0.71 <sub>b2</sub>	7.00±2.83 <sub>b3</sub>
	0.40	15.50±0.71 <sub>a1</sub>	6.00±2.83 <sub>c2</sub>	7.00±4.24 <sub>c3</sub>	7.00±1.41 <sub>b3</sub>
Leucocyte(x104/mm)	Control	2.47±0.08 <sub>a1</sub>	2.50±0.02 <sub>a1</sub>	2.49±0.16 <sub>a1</sub>	2.47±0.04 <sub>a1</sub>
	0.10	2.54±0.04 <sub>b1</sub>	3.38±0.60 <sub>b2</sub>	4.28±0.11 <sub>b3</sub>	4.84±0.01 <sub>b4</sub>
	0.40	2.94±0.10 <sub>b1</sub>	3.96±0.06 <sub>c2</sub>	4.55±0.19 <sub>c3</sub>	5.55±0.04 <sub>c4</sub>

Table (2) : Effect of different lead concentration on the differential white blood cell count of grey mullet Mugil cephalus L

The leucocytes count in the control group did not vary ( $p > 0.05$ ) significantly during the study. The leucocytes counts increased significantly with increasing Pb concentrations and exposure period. Five different subspecies of leucocytes (lymphocytes, monocytes, neutrophils, eosinophils and basophils) were identified in the fish during the study. The lymphocyte and the monocytes constituted the a granulocytes identified while neutrophil, eosinophils and basophils were the granulocytes recorded due to the presence of granules in their cytoplasm. The lymphocytes were the most abundant leucocytes group identified in the blood of the fish exposed to lead acetate. Both small and large lymphocytes were found during the study and they accounted for more than 80 % of the white blood cells. The lymphocytes increased significantly ( $p < 0.05$ ) in the lead-exposed fish when compared with the control group.

The lymphocytes were significantly different ( $p > 0.05$ ) in the treatment groups and the lymphocytosis was both concentration and duration dependent. The monocytes decreased in the lead-exposed fish on the 7th day and thereafter, it increased significantly in the treatment groups ( $p < 0.05$ ) when compared with the control group. The basophiles were the most abundant granular leucocytes in the peripheral blood of Mugil cephalus, L- exposed to lead acetate. The proportion was highest in the first week and on day 21 of the study. The neutrophils are the second largest granular white blood cell in the fish while the eosinophils are the least abundant subpopulation.

#### Effect on serum glucose

The glucose concentration in the control group did not vary throughout the study (Table 3).

Parameters	Concentration(mg/L)	Duration( days)			
		7	14	21	28
Glucose(mg/dl)	Control	17.25± 2.08	17.6±18	17.4±18	
	0.10	26.50± 2.12	32.45± 3.12	41.52± 08.12	52.50± 2.12
	0.40	30.50± 0.70	44.60± 4.12	56.40± 3.12	70.00± 2.83
Cholesterol(mg/dl)	Control	54.06±4.50	55.06±6.40	56.06±3.52	57.05±5.40
	0.10	122.5±3.53	156.5±4.63	187.4±2.50	214.0± 1.80
	0.40	152.5±3.54	197.5±2.51	243.5±3.53	308± 5.19
Total Protein(g/dl)	Control	3.9±05	4.1±03	4.1±0.03	4.1±01
	0.10	4.06±0.06	17.6±18	4.72±0.06	5.28±0.05
	0.40	4.54±0.37	32.45± 3.12	5.14±0.06	38±0.19

**Table (3) :** change in some Biochemical parameters in grey mulle Mugil cephalus L exposed to different Pb doses for different exposure periods.days.

Value in the same column with the same superscript (lower case) are not significantly different (p = 0.05)between different concen-

trations within the same exposure duration. Values with different numeric superscripts differ significantly (P=0.05) between different exposure periods within the same concentration.

**Discussion:**

The results of the study show that lead has significant effects on the leucocyte count of Mugil cephalus L- and the small lymphocytes decreased in number while the neutrophils increased. On the contrary, [20] found decreased leucocyte counts in Clarias gariepinus exposed to 0.45mg Pb /L for 4 to 5 days. The general leucocytosis reported in this study is consistent with the observation of [21] in the dogfish exposed to 50µg/L Cd for 4 days.

Changes of blood glucose are a good indicator of metal stress in fish [22] and alterations in the glucose level might be related to renal injury, liver damage, and lack of nutrition [23]. This study showed a dose- dependent increase of glucose level after first 15-days. The observed increase in the serum glucose level in Mugil Cephalus L, exposed to lead in this study is in accord with report of some earlier workers. Similar increase in plasma glucose was also reported in Prochidolus lineatus exposed to lead [24] and in Oreochromis niloticus exposed to copper [25]. It has been widely reported that hyperglycaemia in fish arises due to the stimulation of catecholamines and corticosteroids [26] .

The increased serum glucose level in this study is an evidence of stress due to lead exposure as [27] argued that coping with such stress is an energy demanding process that requires the fish to mo-

bilize metabolically energy substrates through intense gluconeogenesis. Glucose being one of such known substrates is mobilized through gluconeogenesis to meet this challenge [28] . Heavy metals have been reported to act antagonistically with glucocorticoids by inhibiting the receptors thereby disrupting the osmotic and mineral regulatory mechanisms [29]. Serum increase in glucose due to toxicants has have been associated with hypothalamus-sympathetic-chromaffin cells [30] instead of the hypothalamus-pituitary-internal axis [31] that is known to have tremendous influence on carbohydrate metabolism.

The results of this study showed that the serum total protein increased significantly in Mugil cephalus, L ,exposed to lead. Similar increases in the serum protein level were reported in Oreochromis niloticus exposed to metals [27]. Also increased plasma protein was reported in Mugil exposed to 0.5ppm copper and cadmium [32] . On the contrary, decreased tissue protein was reported in Oreochromis niloticus treated with cadmium [33] and in Cyprinus carpio when exposed to heavy metals [34] .

The enhanced serum cholesterol in this study is an indication of hypercholesteremia in the fish due to the stimulatory effect on the cholesterol biosynthetic pathway. Reduced serum cholesterol level has been reported in Oreochromis niloticus exposed to some heavy

metals [35], [27] and in *Lepomis macrochirus* exposed to methyl mercuric chloride [36]. Cholesterol concentration in the serum of cadmium exposed fish also showed a different pattern. Another study showed a reduction in cholesterol within 15-days, possibly due to tissue damage in the kidney [37]. On the contrary, in *Oreochromis niloticus*, an increase in cholesterol was seen during a 21 day period due to cadmium [38].

This alteration in cholesterol concentration could be due to hazardous effects of metals on cell membrane. The observed elevated cholesterol in this study could have resulted in part to the adverse effect of lead on the liver leading to altered cholesterol metabolism resulting in increased serum cholesterol. This according to [38] could be due to liver and kidney failure that resulted in the release of cholesterol into the blood stream. Thus, increase in cholesterol levels are good indicators of environmental stress in fishes.

### Conclusion:

Generally, the leucocytosis observed in this study gives indication that exposing the fish to lead predisposes it to secondary infections. Also, the reported hyperglycemia, increased serum protein and cholesterol levels are indications of altered carbohydrate, lipid and protein metabolism in the fish due to lead exposure. Serum parameters which provide information as to state of the internal environment of the fish are known to respond quickly to changes in the water quality. The changes in these biomarkers are a reflection of organ dysfunction in the fish due to metal exposure and these biomarkers could be used in ecotoxicological assessment and as early warning indicators of pollution.

### Reference:

1. Jackson RN, Baird D and Els S (2005). [The effect of the heavy metals lead and zinc on the brood and larval development of the burrowing crustacean, \*Callinectes krassii\*](#). *Water S. A* 31: 107-116
2. TNO Report (2001). [Risk to health and the environment related to the use of lead in products](#). TNO Strategy, Technology and Policy. STB-01-39
3. ATSDR (1999). Toxicological profile for lead (Update). Prepared by Research Triangle institute.
4. Jezierska B, Witeska M (2006). [The metal uptake and accumulation in fish living in polluted waters](#). NATO Science Series, Netherlands: Springer
5. Castro-González MI, Méndez-Armenta M (2008). [Heavy metals: Implications associated to fish consumption](#). *Environ. Toxicol. Pharmacol.* 26: 263-271.
6. Katti SR, Sathyanesan AG (1983). [Lead nitrate induced changes in lipid and cholesterol levels in the freshwater fish \*Clarias batrachus\*](#). *Toxicol. Lett.* 19: 93-96.
7. Iger Y, Abraham M (1997). [Rodlet cells in the epidermis of fish exposed to stressors](#). *Tissue Cell*, 29: 431-438.
8. Olojo EAA, Olurin KB, Mbaka G, Oluwemimo AD (2005). [Histopathology of the gill and liver tissues of the African catfish, \*Clarias gariepinus\* exposed to lead](#). *Afr. J. Biotechnol.* 4: 117-122.

9. El-Badawi AA (2005). Effect of lead toxicity on some physiological aspects of Nile tilapia fish, *Oreochromis niloticus*. *Inter. Conf. Vet. Res. Div., NRC, Cairo, Egypt.*
10. Delistraty, D. and Stone, A. (2007). [Dioxins, metals, and fish toxicity in ash residue from space heaters burning used motor oil](#). *Chemosphere*, 68(5): 907-914.
11. Dave G, Xiu R (1991). [Toxicity of mercury, copper, nickel, lead, and cobalt to embryos and larvae of zebrafish, \*Brachydanio rerio\*](#). *Arch Environ Contam Toxicol* 21: 126-134.
12. Hou JL, Zhuang P, Zhang LZ, Feng L, Zhang T, Liu JY, et al. (2011) [Morphological deformities and recovery, accumulation and elimination of lead in body tissues of Chinese sturgeon, \*Acipenser sinensis\*, early life stages: a laboratory study](#). *J Appl Ichthyol* 27: 514-519.
13. Finney D. J. (1971). Probit analysis, 3rd edn. Cambridge University Press, Cambridge, pp 25-26.
14. El-Sherif M.S, Ahmed M, El-Danasoury M.A and El-Nwshy N H.K (2009). [Evaluation of diazinon toxicity on Nile tilapia \(\*Oreochromis niloticus\*\)](#). *J. Fish. Aquat. Sc.* 4(4): 169-177
15. Puchkov N (1964). The white blood cell. In: *The Technique For the Investigation of Fish Physiology.* (Edited by Monson S.) pp 11-15 Jerusalem.
16. Gornall, A.G, Bardawill, C.J and David M.N (1949). [Determination of serum protein by the biuret reaction](#). *J. Biol. Chem.* 177: 751-766
17. Roe J.H (1955). [Determination of sugars in the blood and spinal fluid by anthrone reagent](#). *J. Biol. Chem.* 212: 335-343
18. Abell, L.L and Ley B.B (1952). [A simplified method for the estimation of total cholesterol in the serum and demonstration of its specificity](#). *J. Biol. Chem.* 195: 357
19. Steel R.G.D and Torrie J.H (1960). [Principles and procedures of statistics](#). McGraw-Hill Book. New York
20. Svobodova Z, Vykasova B and Machova J (1994). Sublethal chloride effects of pollutants on freshwater fish (Muller R ed) In: Lloyd R Lugano pp 39-52
21. Annune PA and Ahuma F.T.A (1998). Haematological changes in mud catfish, *Clarias gariepinus* (Burch.) exposed to sublethal concentrations of copper and lead. *J. Aquat. Sc.* 13:33-36
22. Tort L and Hernandez-Pascual M.D (1990). [Haematological effects in dogfish \(\*Scyliorhinus canicula\*\) after short term sublethal cadmium exposure](#). *Acta Hydrochim. Hydrbiol.* 18: 379-383
23. Gagnon A, Jumarie C, Hontela A: [Effects of Cu on plasma cortisol and cortisol secretion by adrenocortical cells of rainbow trout \(\*Oncorhynchus mykiss\*\)](#). *Toxicol* 2006, 78:59-65.
24. Pratap HB, Wendelaar Bonga SE (1990). [Effect of waterborne cadmium on plasma cortisol and glucose in the cichlid fish \*Oreochromis mossambicus\*](#). *Comp Biochem Physiol* , 95:313-317.
25. Martinez C.B.R, Nagae M.Y, Zaia C.T.B.V and Zaia D.A.M (2004). [Acute morphological and physiological effects of lead in](#)

- the neotropical fish, *Prochidolus lineatus*. *Braz. J. Biol.* 64: 797-807
26. Monterio, S.M, Mancera J.M, Fernandes A.F and Sousa M (2005). **Copper induced alterations of the biochemical parameters in the gill and plasma of *Oreochromis niloticus***. *Comp. Biochem. Physiol.C* 141: 375- 383
27. Brown J.A (1993). **Endocrine responses to environmental pollutants**. In: Rankin JC and Jesen FB eds. *Fish Ecophysiology*. Chapman and Hull, London UK pp 276-296
28. Özgür F and Kargm F (2010). **Individual and combined effects of heavy metals on serum biochemistry of Nile tilapia *Oreochromis niloticus***. *Arch. Environ. Contam. Toxicol.* 58: 151-157
29. Vosyliene M.Z (1999). **The effect of heavy metals on haematological indices of fish**. *Act. Zool. Lith. Hydro.* 9: 76-82
30. Wendelaar-Bonga S. E. (1997). **The stress response in fish**. *Physiol. Rev.* 7: 591-625
31. McDonald G and Milligan, L (1997). Ionic, osmotic and acid-base regulation in stress. In: Iwama GK, Pickering AD, Sumpter JP and Schreck CB (eds.). *Fish stress and Health in Aquaculture*. Cambridge University Press, Cambridge, pp 5-118
32. Emad H A. El-Naga , El-Moselhy K.M and Mohamed A.H (2005). **Toxicity of cadmium and copper and their effect on serum biochemical parameters of marine fish *Mugil scheli*** *Egyptian J. Aquat. Res.* 31(2): 60:71
33. Arends, R.J, Mancers J.M, Munoz J.L, Wendelaar-Bonga S.E and Flik G (1999). **The stress response of the gill of seabream (*Sparus aurata* L.) to air exposure and confinement** *J. Endocrinol.* 163: 149-157
34. Almeida JA, Novelli, ELB, Dal-Pai Silva M and Alves-Junior R (2001). **Environmental cadmium exposure and metabolic responses of the Nile tilapia *Oreochromis niloticus*** *Environ. Pollut.* 114:169-175
35. Ruparella, S G; Werma, Y; Saiyed S.R and Rawal U.M (1990). **Effect of cadmium on blood of tilapia, *Oreochromis mossambicus* ( Peters) during prolonged exposure**. *Bull. Environ. Contam. Toxicol.* 45: 305-312
36. Dutta, H. M. and Haghghi, A Z. (1986). **Methylmercuric chloride and serum cholesterol levels in bluegill *Lepomis macrochirus***. *Bull. Environ. Contam. Toxicol.* 36: 181-5.
37. M Saeed Heydarnejad1, Mozhdeh Khosravian-Hemamai1 and Amin Nematollahi(2013). **Effects of cadmium at sub-lethal concentration on growth and biochemical parameters in rainbow trout (*Oncorhynchus mykiss*)** Heydarnejad et al. *Irish Veterinary Journal*, 66:11
38. Oner M, Atli G, Canli M(2008) . **Changes in serum parameters of freshwater fish *Oreochromis niloticus* following prolonged metal (Ag, Cd, Cr, Cu, Zn) exposures**. *Environ Toxicol Chem*, 2:360–366.