

Evidence of Fab Fragment Gene in an Invertebrate : The Sea Star Asterias

Michel Leclerc^{*1}, Nicolas Kresdorn²

¹556 rue Isabelle Romée Sandillon, France

²GenXpro de , Frankfurt, Germany

Corresponding Author: Michel Leclerc, 556 rue Isabelle Romée Sandillon, France. E-mail : mleclerc45@gmail.com

Citation: Michel Leclerc et al. (2016), Evidence of Fab Fragment Gene in an Invertebrate : The Sea Star Asterias Rubens. Int J Biotech & Bioeng. 2:1, 37-38. DOI:[10.25141/2475-3432-2016-1.0052](https://doi.org/10.25141/2475-3432-2016-1.0052)

Copyright: ©2016 Michel Leclerc et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Received: August 23, 2016; **Accepted:** September 19, 2016; **Published:** September 29, 2016

Abstract:

Fc receptor gene, was just discovered with the emergence of Fab fragments gene, in Asterias rubens These last ones were described in Ig light chain and Ig heavy chain , when compared to mouse genome.

Introduction:

The purpose of this article is to draw attention to the mass of information that has accumulated on the primitive sea star antibody since about 1983. This work has emanated from our laboratory, those of Gothenburg (Dr Sam Dupont), those of GenX pro (Frankfurt), those of FASTERIS (Switzerland).

Our purpose will be to recall, the homologies between the primitive antibody and the mouse Igkappa chain [1], the presence of two Immunoglobulin sites in this last one, the specificity of this « antibody » to the antigen HRP which induces the reaction : »antigen-antibody«, at last the existence in the sea star immune system of the Fc receptor gene [2] We present now, the last results, concerning this primitive antibody : the emergence of Fab fragments in the immunized and non-immunized sea star transcriptomes, when compared to mouse genome.

Materials and Methods:

Sea stars were obtained from the Biology Institute (Gothenburg University) Immunizations to HRP (Horse-radish peroxidase), genomic studies were already described [1]. Briefly, after ligation of adapters for Illumina's GSII sequencing system, the cDNA was sequenced on the Illumina GSII platform sequencing. 1.100 bp from one side of the approximately 200 bp fragments sequences were assembled using Velvet (3)

Results:

First a comparison between non-immunized sea star genome and mouse genome was performed. Studies were compared to mouse Immunoglobulin Light chain, then to heavy chain :

One contig (Contig9579) could be annotated via TBLASTX to *Mus musculus* "Mus musculus immunoglobulin IgG2a light chain Fab fragment (Mab10CK) mRNA, partial cds" from the NCBI database (gi|2852986|gb|AF042656.1|), with an e-value of 4.15e-05. On an aligned region of 21 amino acids, 14 positive and 9 identical amino acids were found.

```
5'CAGAGTCACCAGGTGCTGTGTTGACATCAG-  
CACTTTCTGAATCCACCTCCATTGGTTGAT  
CCACTTCCTCTTTCACTATAGTTGACGATTT-  
GATATCAACACTGTCACCTGTAAGCGCAT  
CAACACCGCTTGTTATATCAACAGAC-  
GATCTCCTTTGTGGATCCACTTTTTGATCGGTGG  
ATAGAGGCGGAGTGGATGACTCCATCTTTGCTA-  
ATGTTTTGTCATCGGCTAATGTTGATG  
AACTTTCTTCAGCTTTTACCTCGGTCTTCA-  
CATCCACTGTTTTGTTTTCTGATTTACGG  
AAACAGTTGATGTCGTTTTGACTGGATC-  
CACTGACTTATCAGCCAATGTTCCCTGCGGAG  
TTGTTGATGTACTTTGTTGAGAGATGGCTGT-  
GAAGGGATTTGTGGCGGGATCGAAGGAAG
```

CTCCTAAGCCTTCCCTCCCTGGGGTTGGC-
GTTTCTAGAAGCGAGGAATCTTTGGCATCTT
CGAAACACAAGGGATGGTATGTCTGGAGTT-
GAACCTGACAGCATCTCTCAGTTTCCAAA
CCTCTTCATCTTCATCAAAGAATTGCTCAAAC-
GGCTCATGACACAGAGAACACATCTCAT
CGCCAGTGACGTTTCCATCAATAGCCACA-
CAGAGGATGCCATCTTTATCAGGTTTAGACT
CTGCA G T C A T C T C C T C A A G A A A C T -
GCTCCTCTTTCATCCACTTCGTTGATTTCTTCAA
AGTCCAACCACTGATCGACTTCATAGAAC-
CACTTTCCTGGACTGAACAGCCTTCACTCCAT
CCTTCTCACGACGATTCTGACGGAAGTGC-
CAGTCCAGATGTTTCCGATATCTGTCCATTT
CATCAATGACAAAGCGCATACCA3'

Second, results with mouse Ig heavy chain were obtained, when compared to mouse genome : One contig (c37102_g1_i1) could be annotated via TBLASTX to *Mus musculus* "Mus musculus immunoglobulin IgG2a heavy chain Fab fragment (Mab10CH) mRNA, partial cds" from the NCBI database (gi|2852988|gb|AF042657.1|), with an e-value of 6.1e-06. On an aligned region of 20 amino acids, 14 positive and 10 identical amino acids were found.

5'AACTGCGCCACCAGCAAGTATGCCGTAC-
CCTTCGGCTATTGCGACGGCACCAGACTACCC
CAAAGCAATGGCAGCATCGGCCCTCCATTTG-
GTGGTGAAGCTCCGCCCCCAAGTTACGA
GGAGGCTGTTGGCAAACAGACAGACTGG-
CCAACGCAAGGAACGATGAGGATTACGTCAT
TGACGATCAGTTCGTGCCTAGTTATCCG-
TACTTCAACCTTTCGGTTGATGTACCACGGG
TTCCCGCCATTAACCCCATGGAGTTAAAGAGA3'

At last, results with immunized sea star genome were given when compared to mouse IgG2a light chain : One contig (Contig5278|m.6828) could be annotated via TBLASTX to *Mus musculus* "Mus musculus immunoglobulin IgG2a light chain Fab fragment (Mab10CK) mRNA, partial cds" from the NCBI database (gi|2852986|gb|AF042656.1|), with an e-value of 1.62e-05. On an aligned region of 38 amino acids, 23 positive and 14 identical amino acids were found.

5'TCTTTGGAAACCCCGAACCGGAACTGACCT-
GGCAGAGGAAGAGCCCAGCAGGTGCAGACC
TACCACTCGGTGCAGAGCTCCGATTCAA-
CAACCAGCATTGTATCCTCAGGAATATACAGA
TTGAAGATGCTGGGGAGTATGAATGTATGAC-
CACTAATGGTGGAGGAAGAACATCTGGAA
CCATTACTGTCCGAGTCCCACCCACCTGGCT-
GCAAGAAATCTCTGACATCGAGGGGAACA
TCTACGGAGACTTATCCATGAAGTGTAAG-
CAGAGAGTCCCGATGGCGTAACTACACAT
GGTACCGCAATGCTGAGGTGATCCAAGACCTTC-

CCAGGCATCAGTTTCAGCAACGGTTTGA
AGACACTCACCATCCAGAATCTGACTGTG-
GAAGACACCGGGATGTACCAGTGTATGGCTA
GTAATGAATACGGCCAGATATACACCACAG-
CACAGGTCACCTGTACAAGCCATTGCACCAT
GGTTTGGAGACGCCCTTGAGGTGGACCAGC-
CAGCACCCCGTGGTGGCTCAGTGACCATCC
TCTGCCAACCTAAAGCTGCACCCACACCCGT-
CATCGTCTGGACCAAAGGAAGCGAGACCA
TTGAGAACGGTGGCCGCTACACCGTCATG-
GACAACGGTGATTTGATGATCAGTGGTGTTA
GTGATGCAGATGGAGGAGACTACACCTG-
TACGGCTACCAATAGCATGGGAACTGATTCAA
GCACTGGTAGCTTGATCATCAAGGATGGTA-
CAGCCATGGTGGAGCCACCCATGGCATTGA
CTGTAAACGAAGGGGAGATGGGCACATT-
GACCTGCCGTGCATCACACGATCCGTTCTTTG
AGCTGGAGTATGTGTGGCTGAAAGAAAG-
GAATCCAAGTTGATCCTCGTCTCCCTAACTATG
AGATACCTCAGATCCAAATGGGCGAGCCAAGT-
CAGCTGCATATCAAAGACGCCACTATGG
ATCTAGGAGGGAGTACATGTGC3'

Discussion and Conclusion:

It is clear that mouse Fab IgG2a light chain gene, mouse Fab IgG2a heavy chain gene, are present in the sea star genome. It is also noticeable that these genes, show a « symmetry », or at least a similarity in their configurations : we count on an aligned region of 21 amino-acids, 14 positive and 9 identical in the case of IgG2a light chain, 14 positive and 10 identical in the case of IgG2a heavy chain out of 20 amino-acids.

As for immunized sea star IgG2a light chain which is only studied, in the present work : the number of positive amino-acids increases, in a significant manner (on an aligned region of 38 amino-acids, 23 positive instead 9 in non-immunized sea stars to HRP : undoubtedly it is due to immunization(further studies are bound to unravel the phenomenon). In addition to the work on Fc receptor gene [2], it appears that Fab fragments gene, corroborate the existence of the Invertebrate Primitive Antibody through the *Asterias rubens* history [4].

References:

1. Vincent N, et al, [Metagene 2\(2014\) 320-322](#)
2. Leclerc M., et al [Immunol. Lett \(2016\) submitted](#)
3. Zerbino DR., et al, [Gen Res \(2007\) 18:821-9](#)
4. Leclerc M, « Sea star Genomic studies Following 2 » (2016) Lambert Ed. (Germany)