



## Lactobacillus Rhamnosus BMX 54 Vaginal application usefulness as adjuvant therapy for Preterm Premature Rupture of Membranes (PPROM) in the second Trimester of Pregnancy

Giuseppe Luzi<sup>1\*</sup>, Elisa Iazzetta<sup>2</sup>, Lorenzo Cecconi<sup>1,2</sup>, Silvia Famiani<sup>1</sup>, Saverio Arena<sup>1</sup>, Giancarlo Di Renzo<sup>2</sup> and Giorgio Epicoco<sup>1</sup>

<sup>1</sup>SS CC di Ostetricia e Ginecologia Ospedaliera, Azienda Ospedaliera- Universitaria di Perugia, Italy

<sup>2</sup> Clinica Ostetrico Ginecologica, Azienda Ospedaliera- Universitaria di Perugia, Italy

### Abstract

**Objectives:** Very Preterm Premature Rupture of Membranes (PPROM) complicates about 2% - 5% of pregnancies and leads to an exponential increase in perinatal mortality and morbidity. A lot of Bacteria have been associated with PPRM. The aim of the following study was to evaluate the impact of Lactobacillus BMX 54 vaginal application added to standard of care therapy on pregnancy and neonatal outcome.

**Material and Methods:** This observational study includes 38 consecutive women with PPRM hospitalized in the Department of Obstetric and Gynecology of Perugia Hospital. All the women presented PPRM before of 30 week of pregnancy and have been treated with antibiotics, tocolitics therapy and bed rest: 58% of the pregnancies (22 women) added to standard of care once day vaginal Lactobacillus rhamnosus BMX 54 application while the others (42% of the pregnancies: 16 women) have been treated only with standard therapy without probiotic vaginal application

**Results:** The women who added to standard therapy Lactobacillus rhamnosus BMX 54 daily vaginal application from the beginning of PPRM up to the delivery shown a statistical significant increase ( $p = 0.04$ ) of delivery gestational age when compared with the women treated only with standard therapy. The time course between PPRM and delivery was 26.5 +/- 27.4 days in the Lactobacillus treated group and 11.6 +/- 14.2 days in the other group ( $p = 0.03$ ). CRP (C-Reactive Protein) value at delivery time was 2.37 +/- 2.2 in the lactobacillus treated group and 3.87 +/- 2.5 in the control group ( $p = 0.0003$ ). 5th minute Neonatal APGAR Score was 8.8 +/- 0.8 in treated group and 6.6 +/- 3.7 in control group ( $p = 0.006$ ) while the Hospitalization time of the newborn in Neonatal Intensive Unit Care (NICU) was 43.4 +/- 24 days in the treated group versus 61 +/- 33 days in control group ( $p = 0.01$ )

**Conclusions:** Lactobacillus rhamnosus BMX 54 vaginal application used as adjuvant therapy to standard treatment seems to be a useful approach to reduce PPRM complications improving neonatal and maternal outcomes

**Keywords:** PPRM (Preterm Premature Rupture of Membranes), Lactobacillus Rhamnosus BMX 54, B.V. (Bacterial Vaginosis)

**Corresponding author:** Giuseppe Luzi,

SS CC di Ostetricia e Ginecologia Ospedaliera, Azienda Ospedaliera-Universitaria di Perugia, Italy

E-Mail: giuseppe.luzi1961@libero.it

**Citation:** Giuseppe Luzi et al. (2018). Lactobacillus Rhamnosus BMX 54 Vaginal application usefulness as adjuvant therapy for Preterm Premature Rupture of Membranes (PPROM) in the second Trimester of Pregnancy. *Int J Ped & Neo Heal.* 2:1, 15-20

**Copyright:** © 2018 Giuseppe Luzi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### About Biocore Group

Biocore is an Open Access Journal Publisher and International scientific event organizer which mainly focuses in all the elements of life sciences, medical research, and technology. Biocore is formed solely for exploring the scientific community with a good understanding of the different scientific achievements with a single click from any part of the world. Biocore committed to make genuine and reliable contributions to the scientific community by launching high quality and highly informative Scientific Journals. Our Scientific Conferences gives a path breaking strategies in the modern science and technology.

It is our motto to develop an online scientific platform with Peer Reviewed Scholarly Journals and Scientific Conferences with an esteemed Scientific Alliance by encouraging New Initiatives.

**Received:** December 18, 2017,

**Accepted:** December 28, 2017,

**Published:** January 17, 2018

## Introduction

Preterm Premature Rupture Of Membrane (PPROM) complicates only 2% of pregnancies but is associated with 40% of preterm deliveries and can result in significant neonatal morbidity and mortality <sup>(1-3)</sup>

PPROM is associated with lower latency period from membrane rupture to deliver and during this latency period the ascent of pathogen bacteria from the lower urogenital area could create complications such as intrauterine infections <sup>(4-8)</sup>

Ureaplasma urealyticum, Sneathia species, Escherichia coli, Chlamydia trachomatis, Mycoplasma hominis, Enterococcus faecalis and Gardnerella vaginalis are the common bacteria associated with PPRM <sup>(9)</sup>. Surprising mostly of these bacteria are responsible of Bacterial Vaginosis, the most common infective vaginal pathology of the women all over the world in which the absence of the predominant-lactobacillus species is replaced by gram- bacteria in the vagina <sup>(10)</sup>.

The bacteria spread rapidly after the PPRM has occurred and colonize the surfaces of the amniotic membranes, the umbilical cord and the fetus <sup>(11,12)</sup>.

Low transplacental transfer of antibiotics is responsible for the moderate success of systemic antibiotic therapy <sup>(11,12)</sup>.

Some Investigators have demonstrated that antibiotics could not eliminate the amniotic infection in 83% of PPRM cases <sup>(13)</sup>.

Romero et Coll. <sup>(14)</sup> recently demonstrated that upon clinical admission, 37.3% of preterm PPRM patients delivered within 48 hours of membrane rupture and that only 32.2% of all PPRM patients had a latent phase of more than 14 days.

The three causes of neonatal death associated with PPRM are prematurity, sepsis and pulmonary hypoplasia and women with intrauterine infections deliver earlier than non-infected women and infant born with sepsis have a four times highest mortality than those without sepsis <sup>(15)</sup>

Concluding, there are evidences demonstrating an association between ascending infection from the lower uro-genital tract and PPRM <sup>(16,17)</sup>: in patients with PPRM about one-third of pregnancies have positive amniotic fluid cultures <sup>(16,17)</sup> and studies have shown that bacteria possess the ability to cross intact membranes <sup>(15-17)</sup>.

Standard of care of PPRM, broad-spectrum antibiotics and antenatal corticosteroids, are routinely used in this condition, with very limited success to prevent bacteremia, chorioamnionitis, funisitis and intra-amniotic infection syndrome <sup>(18)</sup>

Taking into account the role of bacteria in PPRM complications and neonatal outcomes and the partial ineffectiveness of systemic antibiotics therapy we thought to add to standard therapy (therapeutic tocolysis, systemic antibiotic and bed rest) in PPRM women a cycle of Lactobacillus rhamnosus BMX 54 (NORMOGIN) vaginal tablet application ( 1 tab/day from PPRM to the delivery) with the aim to

control bacterial replication locally and to improve the fetal/maternal outcomes.

We have selected Lactobacillus rhamnosus BMX 54 between all the available vaginal probiotics because of its effectiveness demonstrated in a large sample size of women (more than 700) affected by Bacterial Vaginosis, in controlled and uncontrolled published clinical trials <sup>(19-24)</sup>.

## Materials and Methods

38 consecutive pregnancy women affected by PPRM before 30 weeks of gestational age hospitalized in the Obstetric and Gynecology department of Perugia University and Hospital department of obstetric and gynecology between 2014 and 2015 entered in this observational trial.

The aim of this observational retrospective study was to preliminary understand if a daily vaginal tablet application of lactobacillus rhamnosus BMX 54 (NORMOGIN) was able to improve pregnancy's delivery time and newborn's score in women affected by PPRM.

All the women have been treated with PPRM standard of care therapy (antibiotics (macrolides or B-lactamic) plus Tocolytics (Atosiban and Progesterone) and bed rest during hospitalization: in 22 (patients of hospital department) of 38 PPRM pregnancies a vaginal tablet containing Lactobacillus rhamnosus BMX 54 (NORMOGIN) has been added daily to the standard therapy.

Medium Gestational Age (GA) together with cervicometria, leukocytes count and CRP at the PPRM (hospitalization time) was collected for every woman.

During hospitalization time in every woman has been collected at alternate days a blood sample for the leukocytes and CRP (C-Reactive Protein) detection and a medium value of every week for these values have been reported.

Delivery time, CRP and leukocytes have also been collected at delivery. In every newborn fetal blood pH, BE (Basis Excess) and 1 minutes and 5 minutes APGAR score have been detected together with number of neonatal deaths, number of neonatal infections and serious neonatal complications.

Data analysis was performed with the help of the SPSS software version 17.0. Average measurements and standard deviation with their respective C I95% were calculated: Student's t test was used for independent data averages for the two groups. Analyses with p less than 0.05 were considered statistically significant.

## Results

Gestational age at PPRM was homogeneous in the two different observational groups (26.19 +/- 2.6 weeks for women treated with lactobacilli too (group A) versus 26 .2 +/- 2.2 weeks for non lactobacillus treated women (group B)) with no statistically significant differences between group A and Group B (p = 0.495)(table 1).

**Tab. 1:** Differences in Gestational Age (G.A.), Cervicometria (Cx), Leukocytes counts (Leuk) and C-Reactive Protein (CRP) at PPRM in the two different groups.

	Group A (Lactobacillus rhamnosus BMX54 group + standard therapy)	Group B (only standard therapy)	Student's T-test for independent data
Number of women	22	16	===
GA at PPRM (weeks)	26.19 ± 2.6	26.2 ± 2.2	p = 0.495 NS
Cx at PPRM (mm)	24.76 ± 11	21.2 ± 9.5	p = 0.16 NS
Leuk at PPRM	11.6 ± 2.8	12.0 ± 4.3	p = 0.355 NS
CRP at PPRM	1.52 ± 2.19	2.04 ± 2.75	p = 0.243 NS

Also cervicometria, leukocytes count and CRP were homogeneous in the two differential observational groups (table 1)

The pregnancy and newborn outcomes has been reported in tab 2; Gestational Age (GA) at delivery time was significant increased in group A patients (29.66 +/- 2.6 weeks) versus group B patients (28 +/- 3 weeks) (p=0.04) and, obviously, also the treatment days after PPRM was significant increased in group A (21 +/- 18 days) versus group B women (11.3 +/- 13.1 days) (p = 0.04) (tab 2).

Also the number of days between PPRM and delivery was significantly higher in group A (26.5 +/- 27.4 days) versus group B (11.6 +/- 14.2) (p=0.03) while newborn outcomes as medium weight at delivery (1299.7 +/- 477 g versus 1161.2 +/- 663 g; p=0.23) and medium weight percentile at delivery (35.7 +/- 19.7 versus 42.3 +/- 15.9%; p=0.14) were not statistically difference between groups (tab 2).

APGAR scores at 1 minute and at 5 minutes were significantly increased in group A newborns versus group B newborns (APGAR 1 min: 6.85 +/- 1.79 versus 5.06 +/- 3.6; p=0.02 and APGAR 5 minutes: 8.8 +/- 0.8 versus 6.6 +/- 3.7; p=0.006); a statistically significative difference between groups were also shown in newborn blood pH (7.2 +/- 0.1 versus 7.07 +/- 0.19; p=0.01), in newborns blood Base Excess (BE: 8.4 +/- 6.6 versus 14.1 +/- 0.1; p = 0.01) and in newborn recovery days in Neonatal Intensive Care Unit (NICU) (43.4 +/- 24 days versus 61 +/- 33 days; p=0.04) (tab 2) Leukocytes counts not significantly differ between groups during the trial observation while CRP was significantly lower in group A women during the second, third and fourth weeks (tab 3)

No newborns was died in lactobacillus treated group (0/22 = 0%) while 4 newborns died in group B women (4/16 = 25%). Also the number of newborn infections (7/22 = 31.8% versus 7/16 = 58.3%) and the number of newborn serious complications (4/22 = 18.1% versus 5/16 = 41.6%) were lower in lactobacillus treated patients versus the control group. No drug's related adverse events have been collected during the trial.

## Discussion

PPROM is associated with lower latency from membrane rupture until

delivery and, consequently, it represents an important cause of perinatal morbidity and mortality<sup>(1-3)</sup>.

During the latency period the ascent of pathogenic microorganisms from the lower uro-genital tract could create complications such as intrauterine infections<sup>(4-8)</sup>.

One of the most common complications in PPRM women is intrauterine infection which could lead to chorioamnionitis, metritis after delivery and perinatal outcome such us neonatal sepsis<sup>(1-8)</sup>.

Perinatal outcomes constitute prematurity, neonatal sepsis, respiratory distress syndrome, intraventricular hemorrhage and risk of fetal and neonatal death<sup>(1-8)</sup>.

Since PPRM causes definite maternal and neonatal morbidity and mortality every effort could be done in order to manage this pathology.

The latency time between PPRM and delivery seems to be a Key point to improve perinatal morbidity and mortality: increase this period could help attending physicians in PPRM management.

Expectant management with antenatal antibiotics and corticosteroids administration are "the recommended standard of care" in the setting of PPRM at gestational age of less than 34 weeks<sup>(18)</sup>.

In this observational retrospective clinical trial we observed that by adding vaginal probiotic (Lactobacillus rhamnosus BMX 54 - NORMOGIN) to the standard of care immediately after PPRM we obtained an increase in latency time between PPRM and delivery improving neonatal outcome.

Taking into account that no differences has been shown between the two groups of women (standard of care plus lactobacillus rhamnosus BMX 54 - group A and only standard of care treatment - group B) at PPRM time in terms of gestational age, CRP, leukocytes values and cervical index it was really surprising to observe a significant increase in latency time between PPRM and delivery only adding vaginal probiotics (table 2).

**Tab. 2:** Results – Pregnancy and newborn outcomes.

	Group A	Group B	Student's T-test for independent data
G.A. at delivery (weeks)	29.66 ± 2.6	28 ± 3	p = 0.04
Treatment days after PPROM (days)	21 ± 18	11.3 ± 13.1	p = 0.04
Days from PPROM after delivery (days)	26.5 ± 27.4	11.6 ± 14.2	p = 0.03
Medium newborn weight at delivery/(gr)	1299.7 ± 477	1161.2 ± 663	p = 0.23
Medium newborn percentile weight at delivery (%)	35.7 ± 19.7	42.3 ± 15.9	p = 0.14
APGAR score 1 minute	6.85 ± 1.79	5.06 ± 3.6	p = 0.03
APGAR score 5 minutes	8.8 ± 0.8	6.6 ± 3.7	p = 0.02
Newborn blood pH	7.2 ± 0.1	7.07 ± 0.19	p = 0.01
Newborn blood Base Excess (BE)	8.4 ± 6.6	14.1 ± 0.1	p = 0.01
Neonatal hospitalization days in NICU (days)	43.4 ± 24	61 ± 33	p = 0.04

More the doubling the days between PPROM and delivery (26.5 days versus 11.6 days:  $p=0.03$ ) obtained only by adding lactobacillus rhamnosus BMX 54 (one vaginal tablet daily) to the PPROM standard of care was an intriguing observation especially if related to the best neonatal outcomes obtained in terms of APGAR Score after 1 and 5 minutes, neonatal blood pH and Base Excess (table 2).

Also the differences in terms of hospitalization days in Neonatal Intensive Care Unit between the two groups of newborn (43.4 versus 61 days:  $p=0.04$ ) seems to encourage the use of vaginal probiotics during PPROM (table 2).

Also relevant was the observation that in group A patients no neonatal death has been detected while 4 newborns was died in group B patients: also neonatal serious complications and neonatal infections were higher in women treated only with standard of care confirming that lactobacillus rhamnosus BMX vaginal addition to the standard of care could improve PPROM outcomes.

It's not surprising that a selected lactobacillus, that has recently demonstrated to be able to counteract Bacterial Vaginosis (BV) recurrences in more than 700 women affected by BV (20-24), could be able to improve neonatal and maternal outcome in women affected by PPROM.

Since bacteria like *Ureaplasma urealyticum*, *Escherichia coli*, *Mycoplasma hominis*, *Enterococcus faecalis* and *Gardnerella vaginalis* are the common bacteria associated with PPROM (9) it could be possible that by restoring vaginal ecosystem we can control gram- vaginal pathogens overgrowth in vagina and their spread up versus amniotic membranes.

CRP movements during the first weeks after PPROM in lactobacillus treated group seems to confort this idea making lactobacillus supplementation "a natural defense" against gram- bacteria overgrowth: we

also have to consider that systemic antibiotic therapy could decrease lactobacillus populations in vagina making the ecosystem more available to faecal and urogenital gram- bacteria overgrowth.

Amniotic infections related to gram- facultative pathogens could represent a "break-down" in PPROM maternal and neonatal complications making lactobacillus rhamnosus BMX 54 vaginal supplementation "a new strategy" to counteract these infections: best APGAR scores at 1 and 5 minutes together with significant differences in neonatal blood pH and Base Excess (BE) seems to correlate with this observation.

On the other hand only *Lactobacillus rhamnosus* has demonstrated to be able to colonize vaginal ecosystem during the 7th weeks after implantation (25) making *Lactobacillus rhamnosus* BMX 54 vaginal administration a reasonable approach to restore vaginal ecosystem.

A significant difference in hospitalization days in NICU (Neonatal Intensive Care Unit) and in neonatal complications also seems to demonstrate that only by restoring vaginal ecosystem by lactobacillus application we can improve neonatal outcome in PPROM women.

Obviously, this retrospective, observational study has a lot of methodological limitations: the study is not a prospective, randomized, double blind-placebo controlled clinical trial; the control group (group B) is "unblinded"; the sample size is too small and the number of patients is different in the two groups, nevertheless, the results obtained only by adding lactobacillus rhamnosus BMX 54 1 vaginal tablet daily to the standard of care are surprising in term of maternal and neonatal outcomes.

If these results will be confirmed in prospective, double blind, placebo-controlled clinical trials vaginal probiotic application could become a new interesting therapeutic strategy to control PPROM when adding to standard of care.

**Tab. 3:** Results – Maternal Leukocites and CRP during the trial

	Group A	Group B	Student's T-test for independent data
Leukocites at 1 <sup>st</sup> week	11.7 ± 3.1	10.1 ± 3.6	p = 0.09
Leukocites at 2 <sup>nd</sup> week	10.6 ± 1.8	9 ± 2	p = 0.24
Leukocites at 3 <sup>rd</sup> week	9.8 ± 2	9 ± 1.5	p = 0.28
Leukocites at 4 <sup>th</sup> week	9.34 ± 3.4	9.25 ± 3.8	p = 0.48
CRP at 1 <sup>st</sup> week	2.03 ± 3.9	2.59 ± 2.2	p = 0.329
CRP at 2 <sup>nd</sup> week	0.92 ± 1.2	2.4 ± 2.2	p = 0.03
CRP at 3 <sup>rd</sup> week	0.56 ± 0.51	1.56 ± 1.6	p = 0.05
CRP at 4 <sup>th</sup> week	0.49 ± 0.43	0.57 ± 0.03	p = 0.03
CRP at Delivery	2.37 ± 2.2	3.87 ± 2.5	p = 0.0003

## References

- Blott M, Greenough A. Neonatal outcome after prolonged rupture of the membranes starting in the second trimester. *Arch. Dis. Child.* 1988; 63 (10 Spec. No): 1146-1150.
- Weissmann-Brenner A, O'Reilly-Green C, Ferber A, Divon MY. Values of amniotic fluid index in cases of preterm premature rupture of membranes. *J. Perinat. Med.* 2009; 37 (3) 232-235. Epub ahead of print.
- Mercer BM. Premature rupture of the membrane. In: Petraglia F, Strauss GF, Gabbe SG, Wises G. *Complicated pregnancy*. 4th ed. London: informa health care; 2007: 713-727.
- Lajos GJ, Passini Junior R, Nomura ML, Amaral E, Pereira BG, Milanez H, et al. Cervical bacterial colonization in women with preterm labor or premature rupture of membranes. *Rev. Bras. Ginecol. Obstet.* 2008; 30 (8): 393-399.
- Yoon BH, Kim YA, Romero R, Kim JC, Park KH, Kim MH, et al. Association of oligohydramnios in women with preterm premature rupture of membranes with an inflammatory response in fetal, amniotic, and maternal compartments. *Am. J. Obstet. Gynecol.* 1999; 181 (4): 784-788.
- Gopalani S, Krohn M, Meyn L, Hitti J, Crombleholme WR. Contemporary management of preterm premature rupture of membranes: determinants of latency and neonatal outcome. *Am. J. Perinatol.* 2004; 21 (4): 183-190.
- Kenyon SL, Taylor DJ, Tarnow-Mordi W; ORACLE Collaborative Group. Broad-spectrum antibiotics for preterm, prelabour rupture of fetal membranes: the ORACLE I randomised trial. *Lancet* 2001; 357 (9261): 979-988.
- Pasquier JC, Picaud JC, Rabilloud M, Claris O, Ecochard R, Moret S, et al. Neonatal outcomes after elective delivery management of preterm premature rupture of the membranes before 34 weeks' gestation (DOMINOS study). *Eur. J. Obstet. Gynecol. Reprod. Biol.* 2009; 143 (1): 18-23.
- Tchirikov M, Zhumadilov Z, Winarno AS, Haase R, Buchmann J. Treatment of preterm premature rupture of membranes with oligo-/anhydramnion colonized by multiresistant bacteria with continuous amnioinfusion and antibiotic administrations through a subcutaneous implanted intrauterine port system: a case report. *Fetal Diagn. Ther.* 2015; oct. 9.
- Truter I, Graz M. Bacterial vaginosis: literature review of treatment options with specific emphasis on non-antibiotic treatment. *Af. J. Pharm. Pharmacol.* 2013; 7 (48): 3060-3067.
- Park HS, Ahn BJ, Jun JK. Placental transfer of clarithromycin in human pregnancies with preterm premature rupture of membranes. *J. Perinat. Med.* 2012; 40: 641-646.
- Heikkinen T, Laine K, Neuvonen PJ, Ekblad U. The transplacental transfer of the macrolide antibiotics erythromycin, roxithromycin and azithromycin. *BJOG*, 2000; 107: 770-775.
- Gomez R, Romero R, Nien JK, Medina L, Carstens M, Kim YM, Espinoza J, Chaiworapongsa T, Gonzalez R, Iams JD, Rojas I. Antibiotic administration to patients with preterm premature rupture of membranes does not eradicate intra-amniotic infection. *J. Matern. Fetal Neonatal Med.* 2007; 20: 167-173.
- Romero R, Miranda J, Chaemsaitong P, Chaiworapongsa T, Kusanovic JP, Dong Z, Ahmed AI, Shaman M, Lannaman K, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ, Yeo L, Kim YM. Sterile and microbial-associated intra-amniotic inflammation in preterm prelabour rupture of membranes. *J. Matern Fetal Neonatal Med.* 2014; 29: 1-16.
- Cotton DB, Hill LM, Strassner HT, Platt LD, Ledger WJ. Use of amniocentesis in preterm gestation with ruptured membranes. *Obstet. Gynecol.* 1984; 63: 38-43.
- Galask RP, Varner MW, Petzold CR, Wilbur SL. Bacterial attachment to the chorionic membranes. *Am. J. Obstet. Gynecol.* 1984; 148: 915-28.
- Gyr TN, Malek A, Matchez-Loic F, Altermatt HJ, Bodmer T, Nicolaidis K, et al. Permeation of human chorioamniotic membranes by *Escherichia coli* in vitro. *Am. J. Obstet. Gynecol.* 1994; 170: 223-7
- Mercer BM. Preterm premature rupture of the membranes. *Obstet. Gynecol.* 2003; 101: 178-193.
- Stojanović N, Plečaš D, Plešinac S. Normal vaginal flora, disorders and application of probiotics in pregnancy. *Arch. Gynecol. Obstet.* 2012; 286 (2): 325-32.
- Rossi A, Rossi T, Bertini M, Caccia G. The use of *Lactobacillus rhamnosus* in the therapy of bacterial vaginosis. Evaluation of clinical efficacy in a population of 40 women treated for 24 months. *Arch. Gynecol. Obstet.* 2010; 281 (6): 1065-1069.
- Marcone V, Rocca G, Lichtner M, Calzolari E. Long-term vaginal administration of *Lactobacillus rhamnosus* as a complementary approach to management of bacterial vaginosis. *Int. J. Gynaecol. Obstet.* 2010; 110 (3): 223-6.

22. Parma M, Dindelli M, Caputo L, Radaelli A, Quaranta L, Candiani M. The role of vaginal lactobacillus rhamnosus (Normogin) in preventing bacterial vaginosis in women with history of recurrences, undergoing surgical menopause: a prospective pilot study. *Eur. Rev. Med. Pharmacol. Sci.* 2013; 17 (10): 1399-403.
23. Bertini M. Is Lactobacillus rhamnosus BMX54 vaginal application a good strategy to counteract bacterial vaginosis recurrences? 18th World Congress on Controversies in Ostetrics, Gynecology & Infertility (COGI) 2014 Monduzzi Editoriale Proceedings.
24. Recine M, Palma E, Domenici L, Giorgini M, Imperiale L, Sassu C, Musella A, Marchetti C, Muzii L, Benedetti Panici C. Restoring vaginal microbiota: biological control of bacterial vaginosis. A prospective case-control study using Lactobacillus rhamnosus BMX 54 as adjuvant treatment against bacterial vaginosis. *Arch. Gynecol. Obstet.* 2015. Epub ahead of print.
25. Reid G, Millsap K, Bruce AW. Implantation of Lactobacillus casei var rhamnosus into vagina. *The Lancet*, 1994; 344 (8931): 1229.